

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

**THIS PAGE BLANK (USPTO)**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07D 401/00</b>		<b>A2</b>	(11) International Publication Number: <b>WO 98/46589</b>
			(43) International Publication Date: 22 October 1998 (22.10.98)
(21) International Application Number: PCT/EP98/02129 (22) International Filing Date: 2 April 1998 (02.04.98) (30) Priority Data: MI97A000867          15 April 1997 (15.04.97)          IT (71) Applicant (for all designated States except US): ANGELINI RICERCHE S.P.A. SOCIETA' CONSORTILE [IT/IT]; Pi- azzale della Stazione, I-00040 S. Palomba (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): ALISI, Alessandra [IT/IT]; Via Nemoense, 188, I-00199 Roma (IT). BRU- FANI, Mario [IT/IT]; Via Aldo Moro, 28, I-00040 Castel Gandolfo (IT). CAZZOLLA, Nicola [IT/IT]; Via Virbio, 56, I-00040 Ariccia (IT). GIANNANGELI, Marilena [IT/IT]; Via Clelia, 88, I-00181 Roma (IT). PINZA, Mario [IT/IT]; Via per Cesano Boscone, 24, I-20094 Corsico (IT). (74) Agents: MARCHI, Massimo et al.; Marchi & Partners s.r.l., Via Pirelli, 19, I-20124 Milano (IT).		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished          upon receipt of that report.</i>	
(54) Title: INDAZOLE AMIDE COMPOUNDS AS SEROTONINERGIC AGENTS			
<div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div>			
(57) Abstract  A compound having general formula (I) wherein R <sub>1</sub> , R <sub>2</sub> , R <sub>3</sub> , R' <sub>3</sub> , R <sub>4</sub> , R <sub>5</sub> and R <sub>6</sub> have the meanings stated in the description, acid addition salts thereof with pharmaceutically acceptable organic and inorganic acids and pharmaceutically acceptable quaternary salts thereof.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	IS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NI	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

"Indazole amide compounds as serotonergic agents"


\*\*\*\*\*

The present invention relates to an indazole amide compound  
possessing a serotonergic action, a method for preparing thereof and  
the pharmaceutical compositions containing the same.

Amongst the numerous known families of serotonin receptors, the  
5HT<sub>4</sub> receptors have only recently been identified in the urinary bladder,  
smooth and cardiac muscle and specific areas of the central nervous  
system. Compounds possessing agonistic, partially agonistic and  
antagonistic actions against such receptors are of potential interest in  
pharmacological treatment of disorders of gastrointestinal motility,  
disorders of the central nervous system, urinary incontinence and cardiac  
arrhythmia. The action of such compounds in fact takes place by  
mimicking or antagonising the ability of serotonin to stimulate intestinal  
motility by activation of the enteric neurons, to modulate important  
cerebral processes such as training, memory and anxiety, to induce  
relaxation of the urinary bladder and to increase frequency of atrial  
contraction.

A family of indazole amide compounds has now been found which  
possess affinity with 5HT<sub>4</sub> receptors and which act as partial agonists or  
antagonists of serotonin.

It is therefore a first object of the present invention to provide an  
indazole amide compound having the general formula:



- 2 -

wherein

R<sub>1</sub> is hydrogen;

R<sub>2</sub> is isopropyl;

R<sub>3</sub> and R'<sub>3</sub> are hydrogen;

5 R<sub>4</sub> and R<sub>5</sub> are hydrogen;

R<sub>6</sub> is selected from the group comprising C<sub>1-3</sub> alkyl, C<sub>3-7</sub> cycloalkyl,

heterocyclic ring having from 5 to 6 members where 1 to 4 members  
are heteroatoms, the same or different from each other, selected from  
the group comprising N, O and S, dimethylamino C<sub>1-3</sub> alkyl, methoxy  
10 C<sub>1-3</sub> alkyl, N-phenyl amide, aminosulphonylmethyl, dihydroxy C<sub>2-3</sub>  
alkyl, aryl, aryl substituted by at least a group selected from halogen  
and hydroxy, aryl C<sub>1-3</sub> alkyl;

acid addition salts thereof with pharmaceutically acceptable organic and  
inorganic acids and pharmaceutically acceptable quaternary salts  
15 thereof.

Preferred examples of aryl are phenyl, naphthyl and biphenyl.

Preferred examples of heterocyclic rings are thienyl, furanyl, pyranyl,  
pyrrolyl, imidazolyl, pyrazolyl, isoxazolyl, pyridinyl, pyrazinyl, pyrimidinyl,  
pyridazinyl, furazanyl, pyrrolinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl,  
20 piperidinyl, piperazinyl, morpholinyl, triazinyl, thiazolyl, tetrazolyl and  
thiadiazolyl. Typical examples of R<sub>6</sub> are cyclopropyl, cyclohexyl, pyridinyl,  
tetrazolyl, morpholinyl, methoxymethyl, methoxypropyl, phenyl,  
chlorophenyl, bromophenyl, hydroxyphenyl, phenethyl,  
dimethylaminomethyl, and aminosulphonylmethyl.

25 It is a second object of the present invention to provide a process for  
preparing a compound of the formula (I), acid addition salts thereof with  
pharmaceutically acceptable organic and inorganic acids and  
pharmaceutically acceptable quaternary salts thereof, comprising:

a) acylating a 4-aminomethyl piperidine of the formula:

- 3 -



(II)


wherein

R<sub>3</sub>, R'<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> have the above mentioned meanings, and

5

P is a suitable protecting group;

by means of a 1-alkyl-indazole-3-carboxylic acid halide of the formula:



(III)


10

wherein

R<sub>1</sub> and R<sub>2</sub> have the above mentioned meanings, and

X is halogen,

to give a compound of the formula:



15

(IV)

wherein


R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R'<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and P have the above mentioned meanings,

b) de-protecting a compound of the formula (IV) to give a compound of

20

the formula:


- 4 -



(V)

wherein  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R'_3$ ,  $R_4$  and  $R_5$  have the above mentioned meanings,

- 5 c) alkylating a compound of the formula (V) with a compound of the formula (VI) to give a compound of the formula (I) according to the following reaction scheme:



wherein

$R_1$ ,  $R_2$ ,  $R_3$ ,  $R'_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  have the above mentioned meanings,  
and

Y is halogen,

- 15 d) optionally forming an acid addition salt of an indazole amide compound of the formula (I) with a pharmaceutically acceptable organic or inorganic acid, or a pharmaceutically acceptable quaternary salt of an indazole amide compound of the formula (I).

Typical examples of protecting groups (P) are benzyloxycarbonyl,  
20 benzyl, tertbutoxycarbonyl and trimethylsilylethoxycarbonyl.

Step a) is preferably carried out by reacting a compound of the formula (II) with a compound of the formula (III) in which X is chlorine, in



the presence of a suitable diluent and at a temperature of from 0 to 140°C for a period of time of from 0.5 to 20 hours.

Preferably the diluent is aprotic, polar or apolar. Still more preferably, it is aprotic apolar. Examples of suitable aprotic apolar diluents are aromatic hydrocarbons such as, for example, benzene, toluene and xylenes. Examples of suitable aprotic polar diluents are dimethylformamide and dimethylsulphoxide.

Still more preferably, the reaction is performed at a temperature of from 15 to 40°C for a period of time of from 1 to 14 hours.

In turn, step (b) is carried out according to techniques known to the person skilled in the art of the protecting group (Theodora W. Greene and Peter G.M. Wuts, "Protective groups in organic synthesis", pp. 309-406, John Wiley & Sons, Inc., N.Y., 1991). In the case of benzyl and benzyloxycarbonyl, the deprotection of the protecting group is preferably carried out by catalytic hydrogenation. An example of a suitable catalyst is palladium on activated carbon.

Preferably the deprotection is carried out by hydrogenation in the presence of a suitable diluent such as, for example, a low aliphatic alcohol, a low aliphatic acid and mixtures thereof. An example of a preferred diluent is an ethyl alcohol /acetic acid mixture.

Step c) is preferably performed with a compound of the formula (VI), in which Y is chlorine or bromine in the presence of a suitable acceptor of acids such as, for example, alkali carbonates and bicarbonates, low trialkylamines and a suitable diluent such as, for example, aromatic hydrocarbons, dimethylformamide and aliphatic low alcohols.

Typical examples of preferred organic and inorganic acids for forming addition salts of the present invention (step d) are oxalic, maleic, tartaric, methanesulphonic, sulphuric, phosphoric acid, hydrogen bromide and hydrogen chloride.

- 6 -

Methyl iodide is a typical example of a preferred compound forming a pharmaceutically acceptable quaternary salt of the invention.

5 The preparation of the above mentioned salts comprises addition (step d) of a pharmaceutically acceptable organic or inorganic acid, or of methyl iodide to an indazole amide compound of the formula (I) obtained in step c).

The intermediates of formula (IV) and (V) are new. They are therefore a further object of the present invention.

10 Alternatively, indazole amide compound of the formula (I) can be prepared by acylation of a suitable 4-aminomethyl piperidine with a compound of the formula (III).

15 Typical examples of pathological conditions which might benefit from treatment with a pharmaceutical composition according to this invention are all the pathologies which are responsive to treatment with antagonists of 5-HT<sub>4</sub> receptor such as, for example, gastrointestinal disorders associated with high intestinal motility, such as IBS (irritable bowel syndrome), urinary incontinence, and cardiac arrhythmias such as atrial fibrillation.

20 Preferably, the pharmaceutical compositions of the present invention will be prepared in suitable dosage forms comprising an effective dose of at least one compound of the formula (I) or a pharmaceutically acceptable addition salt thereof or a quaternary salt thereof and at least one pharmaceutically acceptable inert ingredient.

25 Examples of suitable dosage forms are tablets, capsules, coated tablets, granules, solutions and syrups for oral administration; creams, ointments and medicated adhesive strips for topical administration; suppositories for rectal administration and sterile solutions for injectable, aerosol or ophthalmic administration.

30 The dosage forms may also contain other conventional ingredients such as stabilizing agents, preservatives, surfactants, buffers, salts for

- 7 -

adjusting the osmotic pressure, emulsifiers, sweeteners, coloring agents, flavoring agents, and the like.

When required by particular therapies, the pharmaceutical composition of the present invention may contain other  
5 pharmacologically active ingredients whose concomitant administration is therapeutically useful.

The amount of the compound of formula (I) or of a pharmaceutically acceptable salt thereof may vary within a wide range depending on known factors such as, for example, the type of disease to be treated, the  
10 severity of the disease, the patient's body weight, the dosage form, the chosen route of administration, the number of dosage forms administered per day and the effectiveness of the chosen compound of formula (I). However, the optimum amount may be easily and routinely determined by a person skilled in the art.

15 Typically, the amount of the compound of formula (I) or of a salt thereof in the pharmaceutical composition of this invention will be such as to insure an administered dosage level of from 0.001 to 50 mg/kg/day.

The dosage forms of the pharmaceutical composition according to this invention may be prepared according to methods which are known to the  
20 pharmaceutical chemist and comprise mixing, granulation, compression, dissolution, sterilization, and the like.

The following Examples are intended to illustrate the present invention, without limiting it in any way.

#### EXAMPLE 1

##### 25 Preparation of 1-isopropyl-1H-3-indazolecarbonyl chloride

(III:  $R_1 = H$ ,  $R_2 = C_3H_7$ )

##### a) 2-methylpropyl-1-isopropyl-1H-3-indazolecarboxylate

To a solution of 2-methylpropyl-1H-3-indazolecarboxylate ( 50 g; 0.24 moles) in 1,2-dimethoxy-ethane (300 ml) a solution of isopropyl  
30 bromide (27.5 ml; 0.29 moles) in 1,2-dimethoxy-ethane (100 ml) and

- 8 -

KOH (13.5 g; 0.24 moles) was added and the mixture was heated under reflux for 8 hours. After removal of the solvent, the residue was dissolved in toluene (300 ml), the thus obtained solution was washed with 1N NaOH (100 ml), H<sub>2</sub>O (2x100 ml) and then dried and concentrated *in vacuum*. The residue was purified from the isomer 2-methylpropyl-2-isopropyl-2H-3-indazolecarboxylate via flash chromatography (eluent, hexane : ethyl acetate = 95:5) to give the title compound (23 g) as an oil.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, δ): 1.07 (d, J=7Hz, 6H); 1.66 (d, J=7Hz, 6H); 1.95-2.48 (m, 1H); 4.26 (d, J=7Hz, 2H); 4.96 (hept. J=7Hz, 1H); 7.15-7.70 (m, 3H); 8.03-8.33 (m, 1H).

b) 1-isopropyl-1H-3-indazolecarboxylic acid

A suspension of the compound of the Example 1a) (10 g; 0.04 moles) in 0.75N NaOH (100 ml) was heated under reflux for 12 hours. The solution was then cooled, acidified with 6N HCl (40 ml), the solid precipitate was filtered and recrystallized from 1:1 hexane/ethyl acetate to give the title compound (5.5 g), m.p. 162-3° C (Harada H. et al., "Chem. Pharm. Bull.", 43(11), 1912-1930, 1995).

<sup>1</sup>H NMR (DMSO, δ): 1.54 (d, J=7Hz, 6H); 5.13 (hept, J=7Hz, 1H); 7.20-7.65 (m, 2H); 7.85 (d, J=8Hz, 1H); 8.14 (d, J=7Hz, 1H); 13.08 (s broad, 1H).

c) 1-isopropyl-1H-3-indazolecarbonyl chloride

Thionyl chloride (4 ml, 0.054 moles) was added to a stirred solution of the compound of the Example 1b) and the mixture was stirred under reflux for 2 hours. After removal of the solvent *in vacuum*, the residue was recrystallized from hexane to give 3.5 g of the title compound, m.p. 63-4° C.

- 9 -

Elemental analysis for	C	H	N
$C_{11}H_{11}ClN_2O$			
% found:	59.29	5.20	12.76
% calculated:	59.33	4.98	12.58

$^1H$  NMR ( $CDCl_3$ ,  $\delta$ ); 1.69 (d,  $J=7Hz$ , 6H); 5.00 (hept.,  $J=7Hz$ , 1H); 7.20-7.70 (m, 3H); 8.03-8.33 (m, 1H).

## EXAMPLE 2

Preparation of N3-{[1-(2-phenylethyl)-4-piperidiny]methyl}-1-isopropyl-1H-3-indazolecarboxamide hydrochloride (AFR 306)

(I:  $R_1 = R_3 = R'_3 = R_4 = R_5 = H$ ,  $R_2 = C_3H_7$ ,  $R_6 = C_6H_5$ )

[1-(2-phenylethyl)-1-piperidiny]methylamine (3 g; 0.014 moles), prepared as described in EP-A-0 343 307, in toluene (30 ml) was dropped into a suspension of the compound of the Example 1c) (3 g, 0.014 moles) in toluene (30 ml). After 3 hours at room temperature, the solid was filtered, dissolved in  $H_2O$ , made basic with 6N NaOH solution and extracted with  $CH_2Cl_2$  (2x200 ml). The solvent was removed by evaporation, the residue was purified on  $SiO_2$  column (eluent,  $CHCl_3$ : MeOH = 95 : 5) and transformed into the corresponding hydrochloride.

The obtained product (2 g) melted at 211-212° C.

Elemental analysis for	C	H	N	Cl
$C_{25}H_{33}ClN_4O$				
% found:	68.13	7.52	12.78	8.03
% calculated:	68.09	7.54	12.70	8.04

$^1H$  NMR (DMSO,  $\delta$ ); 1.56 (d,  $J=7Hz$ , 6H); 1.50-2.30 (m, 5H); 2.70-3.90 (m, 10H); 5.10 (hept,  $J=7Hz$ , 1H); 7.05-7.63 (m, 7H); 7.81 (d,  $J=8Hz$ , 1H); 8.21 (d,  $J=8Hz$ , 1H); 8.47 (t,  $J=6Hz$ , 1H); 11.05 (s broad, 1H).

IR (KBr):  $\nu_{CO}$  1652 $cm^{-1}$ .

## EXAMPLE 3

Preparation of N3-{[1-(phenylmethyl)-4-piperidiny]methyl}-1-isopropyl-1H-3-indazolecarboxamide

- 10 -

(IV:  $R_1 = R_3 = R'_3 = R_4 = R_5 = H$ ,  $R_2 = C_3H_7$ ,  $P = -CH_2C_6H_5$ )

To a stirred solution of 1-isopropyl-1H-3-indazolecarbonyl chloride (52 g; 0.234 moles) in toluene (300 ml) it was added dropwise a solution of [1-(phenylmethyl)-4-piperidinyl]methylamine, prepared as described in WO 94/10174, (47.7 g; 0.234 moles) in toluene (200 ml). After 5 hours, the solvent was removed by evaporation under reduced pressure. The reaction mixture was treated with 2N NaOH, extracted with dichloromethane and concentrated *in vacuum*. The solid residue (95 g) was recrystallized from 7:3 hexane/ethyl acetate to afford the title compound as a white solid (45 g), m.p. 72-74° C.

Elemental analysis for	C	H	N
$C_{24}H_{30}N_4O$			
% found:	73.78	7.87	14.35
% calculated:	73.81	7.74	14.35

$^1H$  NMR ( $CDCl_3$ ,  $\delta$ ): 1.59 (d,  $J=7Hz$ , 6H); 1.10-2.25 (m, 7H); 2.80-3.15 (m, 2H); 3.27-3.60 (m, 4H); 4.86 (hept,  $J=7Hz$ , 1H); 7.00-7.60(m, 9H); 8.27-8.52(m, 1H).

IR (KBr):  $\nu_{CO}$  1641 $cm^{-1}$ .

15

## EXAMPLE 4

Preparation of N3-(4-piperidinylmethyl)-1-isopropyl-1H-3-indazolecarboxamide hydrochloride

(V:  $R_1 = R_3 = R'_3 = R_4 = R_5 = H$ ,  $R_2 = C_3H_7$ )

A suspension of the product of the Example 3 (28 g; 0.076 moles) in ethyl alcohol (1500 ml) and glacial acetic acid (66 ml) was hydrogenated over 10% Pd-C (13.4 g) at 35 psi for 24 hours. The mixture was filtered and the filtrate concentrated *in vacuum*. The residue was dissolved in water, treated with 5N NaOH and stirred for 2 hours at room temperature. The solid obtained was filtered off (16.6 g) and converted to the corresponding hydrochloride (9.5 g), m.p. 211-214° C (dec.)

25

- 11 -

Elemental analysis for	C	H	N
$C_{17}H_{25}ClN_4O \cdot 1/2H_2O$			
% found:	58.82	7.68	16.36
% calculated:	59.03	7.58	16.20

$^1H$  NMR (DMSO,  $\delta$ ); 1.55 (d, J=7Hz, 6H); 1.31-2.18 (m, 5H); 2.58-3.64 (m, 7H); 5.09 (hept, J=7Hz, 1H); 7.12-7.60 (m, 2H); 7.80 (d, J=8Hz, 1H); 8.20 (d, J=8Hz, 1H); 8.41 (t, J=6Hz, 1H); 8.82-9.60 (m, 2H).  
IR (KBr):  $\nu_{\infty}$  1658  $cm^{-1}$ .

5

## EXAMPLE 5

Preparation of N3-([1-(4-phenylbutyl)-4-piperidiny]methyl)-1-isopropyl-1H-3-indazolecarboxamide oxalate (AFR603)

(I:  $R_1 = R_3 = R'_3 = R_4 = R_5 = H$ ,  $R_2 = C_3H_7$ ,  $R_6 = -CH_2CH_2C_6H_5$ )

To a stirred suspension of the product of Example 4 as free base (5.27 g; 15.6 mmoles) in ethyl alcohol (20 ml),  $K_2CO_3$  (6.5 g; 50 mmoles) and 4-phenylbromobutane ("Braun", B-44, 2872, 1911) (3.6 g, 17.1 mmoles) were added. The reaction mixture was stirred at reflux for 10 hours. After removal of the solvent, the residue was partitioned between ethyl acetate and 1N HCl. The water phase was made basic with 2N NaOH, extracted with ethyl acetate and concentrated *in vacuum*. The solid was converted to the corresponding oxalate salt (2 g), m.p. 154-155° C.

15

Elemental analysis for	C	H	N
$C_{29}H_{39}N_4O_5 \cdot 1/2H_2O$			
% found:	65.87	7.47	10.62
% calculated:	65.52	7.39	10.54

$^1H$  NMR (DMSO,  $\delta$ ); 1.55 (d, J=7Hz, 6H); 1.31-2.18 (m, 5H); 2.30-3.64 (m, 14H); 5.08 (hept, J=7Hz, 1H); 7.12-7.60 (m, 7H); 7.80 (d, J=8Hz, 1H); 8.19 (d, J=8Hz, 1H); 8.41 (t, J=6Hz, 1H).

20

- 12 -

## EXAMPLE 6

Preparation of N3-([1-(2-cyclohexylethyl)-4-piperidinyl]methyl)-1-isopropyl-1H-3-indazolecarboxamide hydrochloride (AFR604)

(I:  $R_1 = R_3 = R'_3 = R_4 = R_5 = H$ ,  $R_2 = C_3H_7$ ,  $R_6 = C_6H_{11}$ )

- 5 Following the procedure of Example 5, N3-(4-piperidinylmethyl)-1-isopropyl-1H-3-indazolecarboxamide (4.42 g) and (2-bromoethyl)-cyclohexane ("J.A.C.S.", 48, 1089-1093, 1926) (4.63 g) gave the title compound (2.5 g), m.p. 244-246° C (dec.)

Elemental analysis for	C	H	N	Cl <sup>-</sup>
$C_{25}H_{39}N_4O \cdot 1/2 H_2O$				
% found:	65.51	9.05	12.57	7.89
% calculated:	65.83	8.84	12.28	7.77

- 10  $^1H$  NMR (DMSO,  $\delta$ ); 1.55 (d, J=7Hz, 6H); 0.68-2.18 (m, 17H); 2.63-3.70 (m, 10H); 5.09 (hept, J=7Hz, 1H); 7.12-7.60 (m, 2H); 7.80 (d, J=8Hz, 1H); 8.20 (d, J=8Hz, 1H); 8.41 (t, J=6Hz, 1H); 10.70 (s broad 1H).  
IR (KBr):  $\nu_{\infty}$  1656  $cm^{-1}$ .

## EXAMPLE 7

Preparation of N3-([1-[3-(dimethylamino)propyl]-4-piperidinyl]methyl)-1-isopropyl-1H-3-indazolecarboxamide dimaleate (AFR606)

(I:  $R_1 = R_3 = R'_3 = R_4 = R_5 = H$ ,  $R_2 = C_3H_7$ ,  $R_6 = -CH_2NC_2H_5$ )

- 15 Following the procedure of Example 5, N3-(4-piperidinylmethyl)-1-isopropyl-1H-3-indazolecarboxamide (3 g) and N-(3-chloropropyl)-N, N-dimethylamine hydrochloride (580 mg) gave the title compound (950 mg),  
20 m.p. 155-156° C.

Elemental analysis for	C	H	N
$C_{30}H_{43}N_5O_9 \cdot 1/2 H_2O$			
% found:	57.83	7.01	11.11
% calculated:	57.50	7.08	11.18



- 13 -

$^1\text{H}$  NMR (DMSO,  $\delta$ ); 1.55 (d,  $J=7\text{Hz}$ , 6H); 1.68-2.28 (m, 7H); 2.81 (s, 6H); 2.75-3.75 (m, 11H); 5.09 (hept,  $J=7\text{Hz}$ , 1H); 6.09 (s, 4H); 7.12-7.60 (m, 2H); 7.81 (d,  $J=8\text{Hz}$ , 1H); 8.20 (d,  $J=8\text{Hz}$ , 1H); 8.45 (t,  $J=6\text{Hz}$ , 1H).

## EXAMPLE 8

5      Preparation of N3-((1-[2-(4-morpholinyl)ethyl]-4-piperidinyl)methyl)-1-isopropyl-1H-3-indazolecarboxamide dihydrochloride (AFR607)

(I:  $R_1 = R_3 = R'_3 = R_4 = R_5 = \text{H}$ ,  $R_2 = \text{C}_3\text{H}_7$ ,  $R_6 = \text{C}_4\text{H}_4\text{NO}$ )

Following the procedure of Example 5, N3-(4-piperidinylmethyl)-1-isopropyl-1H-3-indazolecarboxamide (3 g) and 4-(2-chloroethyl)-morpholine (3.42 g) gave the title compound (3.2 g), m.p. 266-267° C (dec.)

Elemental analysis for	C	H	N	Cl
$\text{C}_{23}\text{H}_{37}\text{Cl}_2\text{N}_5\text{O}_2 \cdot 1/2 \text{H}_2\text{O}$				
% found:	55.74	7.61	13.96	14.12
% calculated:	55.75	7.73	14.13	14.31

$^1\text{H}$  NMR (DMSO,  $\delta$ ); 1.55 (d,  $J=7\text{Hz}$ , 6H); 1.30-2.25 (m, 5H); 2.75-4.30 (m, 19H); 5.09 (hept,  $J=7\text{Hz}$ , 1H); 7.12-7.60 (m, 2H); 7.81 (d,  $J=8\text{Hz}$ , 1H); 8.20 (d,  $J=8\text{Hz}$ , 1H); 8.45 (t,  $J=6\text{Hz}$ , 1H); 10.80 (s broad, 1H); 10.60 (s broad, 1H).  
IR (KBr):  $\nu_{\text{co}}$  1652  $\text{cm}^{-1}$ .

## EXAMPLE 9

20      Preparation of N3-((1-[2-((methylsulphonyl)amino)ethyl]-4-piperidinyl)methyl)-1-isopropyl-1H-3-indazolecarboxamide hydrochloride (AFR703)

(I:  $R_1 = R_3 = R'_3 = R_4 = R_5 = \text{H}$ ,  $R_2 = \text{C}_3\text{H}_7$ ,  $R_6 = \text{CH}_3\text{SO}_2\text{NH}-$ )

Following the procedure of Example 5, N3-(4-piperidinylmethyl)-1-isopropyl-1H-3-indazolecarboxamide (5 g) N-(2-bromoethyl)-methane sulphonamide (WO 93/18036) (3 g) gave the title compound (1.5 g), m.p. 186-187° C (dec.)

- 14 -

Elemental analysis for	C	H	N	S	Cl <sup>-</sup>
$C_{20}H_{32}ClN_5O_3S$					
% found:	52.15	7.22	15.30	6.98	7.77
% calculated:	52.45	7.04	15.29	7.00	7.74

<sup>1</sup>H NMR (DMSO,  $\delta$ ); 1.55 (d, J=7Hz, 6H); 1.40-2.30 (m, 5H); 3.00 (s, 3H); 2.75-3.80 (m, 10H); 5.09 (hept, J=7Hz, 1H); 7.12-7.70 (m, 3H); 7.80 (d, J=8Hz, 1H); 8.20 (d, J=8Hz, 1H); 8.45 (t, J=6Hz, 1H); 10.73 (s broad, 1H).

5 IR (KBr):  $\nu_{\text{co}}$  1651 cm<sup>-1</sup>.

#### EXAMPLE 10

##### Preparation of N3-({1-[2-(2-pyridinyl)ethyl] 4-piperidinyl)methyl}-1-isopropyl-1H-3-indazolecarboxamide hydrochloride (AFR605)

(I: R<sub>1</sub> = R<sub>3</sub> = R'<sub>3</sub> = R<sub>4</sub> = R<sub>5</sub> = H, R<sub>2</sub> = C<sub>3</sub>H<sub>7</sub>, R<sub>6</sub> = C<sub>5</sub>H<sub>4</sub>N)

10 To a stirred suspension of the product of Example 4 as free base (10 g; 33.3 mmoles), 2-vinylpyridine (3.6 g; 34 mmoles), glacial acetic acid (2 ml) and water (2.5 ml) were added. After 16 hours at 95° C, the reaction mixture was made basic with 2N NaOH, extracted with ethyl acetate and concentrated *in vacuum*. The residue was purified by flash silica-gel

15 chromatography with CHCl<sub>3</sub>:MeOH = 97 : 3 as eluent to yield a solid which was converted to hydrochloride salt (5 g), m.p. 122-123° C (dec.)

Elemental analysis for	C	H	N	Cl <sup>-</sup>
$C_{24}H_{32}ClN_5O \cdot H_2O$				
% found:	62.80	7.42	15.18	7.78
% calculated:	62.66	7.45	15.22	7.71

<sup>1</sup>H NMR (DMSO,  $\delta$ ); 1.55 (d, J=7Hz, 6H); 1.68-2.30 (m, 5H); 2.80-3.78 (m, 12H); 5.10 (hept, J=7Hz, 1H); 7.12-7.60 (m, 4H); 7.68-8.00 (m, 2H); 8.21 (d, J=7Hz, 1H); 8.33-8.70 (m, 2H); 11.05 (s broad, 1H).

20 IR (KBr):  $\nu_{\text{co}}$  1644 cm<sup>-1</sup>.

#### TEST 1

##### Antagonistic Action on 5-HT<sub>4</sub> Receptor

- 15 -

The antagonistic action of the compounds of the formula (I) was evaluated by testing the influence of the compound under evaluation on serotonin-induced relaxation of rat oesophageal tunica pre-contracted with carbachol according to the method described by J.D. Gale et al. in "Br. J. Pharmacol.", 111, 332-338, (1994).

All the tested compounds of the invention showed a  $pA_2 > 8$ . The specific values for AFR 603, AFR 604, AFR 605 and AFR 306 are shown Table 1 below.

Table 1


Compound	$pA_2$	s.e.
AFR 603	9.12	1.42
AFR 604	8.19	0.99
AFR 605	10.8	1.90
AFR 306	9.36	0.38

s.e. = standard error

- 16 -

## CLAIMS

1. A compound having the general formula



(I)

5 wherein:

$R_1$  is hydrogen;

$R_2$  is isopropyl;

$R_3$  and  $R'_3$  are hydrogen;

$R_4$  and  $R_5$  are hydrogen;

- 10  $R_6$  is selected from the group comprising  $C_{1-3}$  alkyl,  $C_{3-7}$  cycloalkyl, heterocyclic ring having from 5 to 6 members where 1 to 4 members are heteroatoms, the same or different from each other, selected from the group comprising N, O and S, dimethylamino  $C_{1-3}$  alkyl, methoxy  $C_{1-3}$  alkyl, N-phenyl amide, aminosulphonylmethyl, dihydroxy  $C_{2-3}$  alkyl, aryl, aryl substituted by at least a group selected from halogen and hydroxy, aryl  $C_{1-3}$  alkyl;


15 acid addition salts thereof with pharmaceutically acceptable organic and inorganic acids and pharmaceutically acceptable quaternary salts thereof.

- 20 2. A compound according to claim 1, characterized in that aryl is selected from the group comprising phenyl, naphthyl and biphenyl.
- 25 3. A compound according to claim 1, characterized in that heterocyclic rings are thienyl, furanyl, pyranyl, pyrrolyl, imidazolyl, pyrazolyl, isoxazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, furazanyl,

pyrrolinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidinyl,  
piperazinyl, morpholinyl, triazinyl, thiazolyl, tetrazolyl and thiadiazolyl.

4. A compound according to claim 1, characterized in that R<sub>6</sub> is  
selected from the group comprising cyclopropyl, cyclohexyl, pyridinyl,  
5 tetrazolyl, morpholinyl, methoxymethyl, methoxypropyl, phenyl,  
chlorophenyl, bromophenyl, hydroxyphenyl, phenethyl,  
dimethylaminomethyl, and aminosulphonylmethyl.
5. A compound according to claim 1, characterized in that R<sub>1</sub>, R<sub>3</sub>, R'<sub>3</sub>,  
R<sub>4</sub> and R<sub>5</sub> are hydrogen, R<sub>2</sub> is isopropyl and R<sub>6</sub> is phenyl.
- 10 6. A compound according to claim 1, characterized in that R<sub>1</sub>, R<sub>3</sub>, R'<sub>3</sub>,  
R<sub>4</sub> and R<sub>5</sub> are hydrogen, R<sub>2</sub> is isopropyl and R<sub>6</sub> is phenethyl.
7. A compound according to claim 1, characterized in that R<sub>1</sub>, R<sub>3</sub>, R'<sub>3</sub>,  
R<sub>4</sub> and R<sub>5</sub> are hydrogen, R<sub>2</sub> is isopropyl and R<sub>6</sub> is cyclohexyl.
8. A compound according to claim 1, characterized in that R<sub>1</sub>, R<sub>3</sub>, R'<sub>3</sub>,  
15 R<sub>4</sub> and R<sub>5</sub> are hydrogen, R<sub>2</sub> is isopropyl and R<sub>6</sub> is pyridinyl.
9. A compound according to claim 1, characterized in that R<sub>1</sub>, R<sub>3</sub>, R'<sub>3</sub>,  
R<sub>4</sub> and R<sub>5</sub> are hydrogen, R<sub>2</sub> is isopropyl and R<sub>6</sub> is  
dimethylaminomethyl.
10. A compound according to claim 1, characterized in that R<sub>1</sub>, R<sub>3</sub>, R'<sub>3</sub>,  
20 R<sub>4</sub> and R<sub>5</sub> are hydrogen, R<sub>2</sub> is isopropyl and R<sub>6</sub> is morpholinyl.
11. A compound according to claim 1, characterized in that R<sub>1</sub>, R<sub>3</sub>, R'<sub>3</sub>,  
R<sub>4</sub> and R<sub>5</sub> are hydrogen, R<sub>2</sub> is isopropyl and R<sub>6</sub> is  
aminosulphonylmethyl.
12. A process for preparing a compound of the formula (I), acid addition  
25 salts thereof with pharmaceutically acceptable organic and inorganic  
acids and pharmaceutically acceptable quaternary salts thereof,  
comprising:  
a) acylating a 4-aminomethyl piperidine of the formula:

- 18 -



(II)


wherein

R<sub>3</sub>, R'<sub>3</sub>, R<sub>4</sub> and R<sub>5</sub> have the above mentioned meanings, and

5

P is a suitable protecting group;

by means of a 1-alkyl-indazole-3-carboxylic acid halide of the formula:



(III)


10

wherein

R<sub>1</sub> and R<sub>2</sub> have the above mentioned meanings, and

X is halogen,

to give a compound of the formula:



15

(IV)


wherein

R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R'<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub> and P have the above mentioned meanings,

b) de-protecting a compound of the formula (IV) to give a compound of the formula:

20


- 19 -



(V)

wherein  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R'_3$ ,  $R_4$  and  $R_5$  have the above mentioned meanings,

- 5 c) alkylating a compound of the formula (V) with a compound of the formula (VI) to give a compound of the formula (I) according to the following reaction scheme:



wherein

$R_1$ ,  $R_2$ ,  $R_3$ ,  $R'_3$ ,  $R_4$ ,  $R_5$  and  $R_6$  have the above mentioned meanings, and


Y is halogen,

- 15 d) optionally forming an acid addition salt of an indazole amide compound of the formula (I) with a pharmaceutically acceptable organic or inorganic acid, or a pharmaceutically acceptable quaternary salt of an indazole amide compound of the formula (I).
13. A process according to claim 12, characterized in that P is selected
- 20 from the group comprising benzyloxycarbonyl, benzyl, tertbutoxycarbonyl, trimethylsilylethoxycarbonyl.
14. A process according to claims 12 or 13, characterized in that step a) is carried out by reacting a compound of the formula (II) with a

- 20 -

compound of the formula (III) in which X is chlorine, in the presence of a diluent and at a temperature of from 0 to 140°C for a period of time of from 0.5 to 20 hours.

15. A process according to claim 13, characterized in that when P is benzyl or benzyloxycarbonyl, step b) is carried out by catalytic hydrogenation.
16. A process according to any of the preceding claims from 12 to 14, characterized in that when, in a compound of the formula (VI), Y is chlorine or bromine, step c) is performed in the presence of an acceptor of acids and in the presence of a diluent.
17. A process according to claim 12, characterized in that methyl iodide forms a pharmaceutically acceptable quaternary salt of a compound of formula (I), step d).
18. An intermediate compound having the general formula:



(IV)

wherein

R<sub>1</sub> is hydrogen;

20 R<sub>2</sub> is isopropyl;

R<sub>3</sub> and R'<sub>3</sub> are hydrogen;

R<sub>4</sub> and R<sub>5</sub> are hydrogen; and


P is a protecting group.

19. A compound according to claim 18, characterized in that P is selected from the group comprising benzyloxycarbonyl, benzyl, 25 tertbutoxycarbonyl, trimethylsilylethoxycarbonyl.



- 21 -

20. A compound according to claim 18, characterized in that  $R_1$ ,  $R_3$ ,  $R'_3$ ,  $R_4$  and  $R_5$  are hydrogen,  $R_2$  is isopropyl and P is benzyl.
21. An intermediate compound having the general formula:



(V)

wherein

$R_1$  is hydrogen;


$R_2$  is isopropyl;

$R_3$  and  $R'_3$  are hydrogen;

$R_4$  and  $R_5$  are hydrogen.

22. A compound according to claim 17, characterized in that  $R_1$ ,  $R_3$ ,  $R'_3$ ,  $R_4$  and  $R_5$  are hydrogen and  $R_2$  is isopropyl.

23. A pharmaceutical composition, characterized in that the said composition comprises an effective dose of at least one compound of the formula:



(I)

wherein

$R_1$  is hydrogen;

- 22 -

R<sub>2</sub> is isopropyl;

R<sub>3</sub> and R'<sub>3</sub> are hydrogen;

R<sub>4</sub> and R<sub>5</sub> are hydrogen;

- 5        R<sub>6</sub> is selected from the group comprising C<sub>1-3</sub> alkyl, C<sub>3-7</sub> cycloalkyl, heterocyclic ring having from 5 to 6 members where 1 to 3 members are heteroatoms, the same or different from each other, selected from the group comprising N, O and S, dimethylamino C<sub>1-3</sub> alkyl, methoxy C<sub>1-3</sub> alkyl, N-phenyl amide, aminosulphonylmethyl, dihydroxy C<sub>2-3</sub> alkyl, aryl, aryl substituted  
10        by at least a group selected from halogen and hydroxy, aryl C<sub>1-3</sub> alkyl;  
acid addition salts thereof with pharmaceutically acceptable organic and inorganic acids and pharmaceutically acceptable quaternary salts thereof.

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : <b>C07D 401/12, A61K 31/415</b></p>	<p><b>A3</b></p>	<p>(11) International Publication Number: <b>WO 98/46589</b> (43) International Publication Date: 22 October 1998 (22.10.98)</p>
<p>(21) International Application Number: PCT/EP98/02129 (22) International Filing Date: 2 April 1998 (02.04.98) (30) Priority Data: MI97A000867 15 April 1997 (15.04.97) IT (71) Applicant (for all designated States except US): ANGELINI RICERCHE S.P.A. SOCIETA' CONSORTILE [IT/IT]; Piazze della Stazione, I-00040 S. Palomba (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): ALISI, Alessandra [IT/IT]; Via Nimorese, 188, I-00199 Roma (IT). BRUFANI, Mario [IT/IT]; Via Aldo Moro, 28, I-00040 Castel Gandolfo (IT). CAZZOLLA, Nicola [IT/IT]; Via Virbio, 56, I-00040 Ariccia (IT). GIANNANGELI, Marilena [IT/IT]; Via Clelia, 88, I-00181 Roma (IT). PINZA, Mario [IT/IT]; Via per Cesano Boscone, 24, I-20094 Corsico (IT). (74) Agents: MARCHI, Massimo et al.; Marchi &amp; Partners s.r.l., Via Pirelli, 19, I-20124 Milano (IT).</p>		<p>(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RJ, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> (88) Date of publication of the international search report: 28 January 1999 (28.01.99)</p>

(54) Title: INDAZOLE AMIDE COMPOUNDS AS SEROTONINERGIC AGENTS



(57) Abstract

A compound having general formula (I) wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>3'</sub>, R<sub>4</sub>, R<sub>5</sub> and R<sub>6</sub> have the meanings stated in the description, acid addition salts thereof with pharmaceutically acceptable organic and inorganic acids and pharmaceutically acceptable quaternary salts thereof.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/02129

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 6 C07D401/12 A61K31/415

According to International Patent Classification(IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 38420 A (NISSHIN FLOUR MILLING CO) 5 December 1996 see abstract see page 64 - page 65; example 7 see page 73; example 18 see page 75; example 21	1-11, 18, 21, 23
P, X	- & EP 0 829 474 A (NISSHIN FLOUR MILLING CO) 18 March 1998 see page 49 - page 50; example 7 see page 56; example 18 see page 57 - page 58; example 21 see claims 1, 10, 16-21 ---	
X	WO 93 03725 A (SMITHKLINE BEECHAM PLC) 4 March 1993 see page 2, line 1 - page 3, line 30 see page 32; example 10 see page 46 - page 47; example 39 ---	1-11, 18, 21, 23
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

8 October 1998

Date of mailing of the international search report

14/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Paisdor, B

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 98/02129

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 732 333 A (LILLY CO ELI) 18 September 1996 see the whole document ---	1,23
A	EP 0 501 322 A (GLAXO GROUP LTD) 2 September 1992 see the whole document -----	1,23

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 98/02129

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9638420 A	05-12-1996	CA 2222532 A EP 0829474 A	05-12-1996 18-03-1998
WO 9303725 A	04-03-1993	AU 2435092 A AU 5194496 A CA 2116024 A EP 0600955 A JP 6510283 T MX 9204786 A NZ 243993 A PT 100785 A ZA 9206208 A AP 373 A AU 668102 B AU 2541892 A AU 691430 B AU 6073596 A BR 9206599 A CA 2118812 A CN 1073173 A CZ 9400560 A EP 0604494 A FI 941178 A WO 9305038 A HU 70154 A JP 6510537 T MX 9205204 A NO 940874 A NZ 244282 A PT 100855 A SG 49153 A SK 30294 A US 5580885 A	16-03-1993 18-07-1996 04-03-1993 15-06-1994 17-11-1994 01-04-1993 26-10-1994 29-04-1994 24-05-1993 07-12-1994 26-04-1996 05-04-1993 14-05-1998 03-10-1996 08-11-1994 18-03-1993 16-06-1993 13-07-1994 06-07-1994 11-03-1994 18-03-1993 28-09-1995 24-11-1994 31-01-1994 11-03-1994 28-08-1995 30-11-1993 18-05-1998 07-12-1994 03-12-1996
EP 0732333 A	18-09-1996	US 5654320 A AU 6894896 A CA 2215359 A CN 1183041 A NO 974221 A PL 322362 A	05-08-1997 18-11-1996 31-10-1996 27-05-1998 27-10-1997 19-01-1998

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/02129

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0732333 A		SK 124297 A	04-02-1998
		WO 9633713 A	31-10-1996
		US 5798367 A	25-08-1998
EP 0501322 A	02-09-1992	AT 142205 T	15-09-1996
		AU 645402 B	13-01-1994
		AU 1209492 A	15-09-1992
		CA 2105179 A	26-08-1992
		DE 69213276 D	10-10-1996
		DE 69213276 T	23-01-1997
		WO 9214727 A	03-09-1992
		ES 2091963 T	16-11-1996
		FI 933697 A	23-08-1993
		HU 65486 A	28-06-1994
		IL 101042 A	18-06-1996
		JP 6505238 T	16-06-1994
		MX 9200762 A	01-08-1992